



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

the mandible : elytra narrower than the prothorax, striate and slightly crenate, especially towards the base, and in a good light.  $8\frac{1}{2}$  lines long.

Var. *a.* Finely and regularly punctate-striate throughout.

Var. *b.* Striæ filled with numerous fine punctures.

*AGRION venerinotata.* ♂ Yellow, eyes reddish brown, connected by a broad black band, a transverse black spot upon the clypeus: thorax black above, with a narrow dorsal line of yellow: abdomen marked above with 3 black ♀-shaped crosses, basal segment dark polished green above, continuing to the middle of the 2d segment, where it ends in a point; wings short. 10 lines long.

*TERMES frontalis.* *Soldier.* Amber color, front deeply notched, a small eye-like tubercle at the inner base of the tentacles: mandibles suddenly incurved, rufo-testaceous at base, changing into black towards the tip; antennæ 15-jointed: legs pale, with the unguis darker. 3 lines long. *Male?* Black, region of the mouth, and legs (except the femora) pale yellowish.  $4\frac{1}{2}$  lines long to the extremity of the wings, which are longer than the body, and extend beyond it more than half their length. Found in or beneath logs, the winged individuals appearing in May in Pennsylvania.

*Correction.*—*Hoplia modesta*, Proceedings, Vol. I., p. 304, is apparently identical with Germar's *Melolontha mucorea*.

### *Meeting for Business, April 30, 1844.*

VICE PRESIDENT MORTON in the Chair.

The Committee, to whom was referred a paper by Dr. Hallowell, describing new species of African reptiles, reported in favour of publication.\*

The Committee on Colonel Abert's communication on a fresh-water univalve, found in the vicinity of Washington, D. C., reported it to be the *Paludina vivipara* of authors.

The Academy then proceeded to ballot for Members and Correspondents, with the following result:

MEMBER.

J. Fisher Leaming, of Philadelphia.

CORRESPONDENT.

Rev. D. Ziegler, of York, Pennsylvania.

\* The publication of this paper is necessarily deferred until the next number.